

Centre de Physique Théorique* - CNRS - Luminy, Case 907 F-13288
Marseille Cedex 9 - France

QUANTUM EFFECTS ASSOCIATED WITH AN ELECTRICALLY NEUTRAL MASSLESS PARTICLE IN A BERTOTTI-ROBINSON UNIVERSE

Víctor M. VILLALBA¹

Abstract

In the present article we obtain, by separation of variables, an exact solution to the Dirac equation with anomalous momentum for an electrically neutral massless particle in a Bertotti-Robinson universe. We discuss the phenomenon of particle creation in this model.

Key-Words : 11.10, 03.65

October 1994
CPT-94/P.3081

anonymous ftp or gopher: cpt.univ-mrs.fr

* Unité Propre de Recherche 7061

¹permanent address: Centro de Física, Instituto Venezolano de Investigaciones Científicas IVIC, Apdo. 21827, Caracas 1020-A Venezuela
e-mail address: villalba@cpt.univ-mrs.fr, villalba@dino.conicit.ve

In the past twenty years there has been a increasing interest in studying quantum effects in cosmological backgrounds. Since the publication of the pioneer works of Parker [1] and Grib *et al* [2] , a great body of articles have been devoted to the problem of studying the creation of scalar and spin 1/2 particles in different cosmological scenarios. In this direction the formulation of a quantum field theory on curved manifolds is required, and consequently we have to solve in the corresponding curved background the relativistic wave functions associated with one-particle states.

Despite the great effort dedicated to the search of exact solutions of relativistic wave equations in curved space-times, the number of exact solvable examples is relatively scarce. This one, in most of the cases, is related to the impossibility of achieving the preliminary step of separating variables in the corresponding wave equations. Therefore we have to restrict ourselves to those metrics permitting such a separation of variables. That is the case for some diagonal metric, as well as the type D Petrov spaces. The presence of conserved quantities as well as a high degree of symmetries of the metric are desirable if we are willing to find exact solution of the Klein-Gordon or the Dirac equations. This is the case for homogeneous spaces like the Friedmann-Robertson-Walker metric.

A good scenario for discussing quantum effects is the Bertotti-Robinson electromagnetic universe [3, 4, 5]. This is a non-asymptotically flat space-time generated by a constant, homogeneous electric field. The metric associated with this universe can be written as

$$ds^2 = \frac{1}{\sqrt{\lambda}} \left[d\vartheta^2 + \sin^2 \vartheta d\varphi^2 + \cos^2 \vartheta dR^2 - dt^2 \right] \quad (1)$$

Among the advantages of the metric (1) we can mention that it can be regarded as a reducible Einstein space , also it has zero Weyl tensor and , because of its diagonal and very simple form, the Klein Gordon and Dirac equations are exactly soluble in terms of special functions[5]. Perhaps the Bertotti Robinson universe is the only model available in the literature where it is possible to solve the wave equations taking into account that the background field is charged and therefore is present the electromagnetic minimal coupling via the vector potential A_μ . Here it is worth mentioning that is not the case for the cosmological models where the electromagnetic interaction goes in the radiation fluid.

Since the metric (1) is conformally flat, creation of electric free massless particles cannot be expected [6]. This situation can be modified if we consider the existence of a no trivial anomalous moment. In this case, there is an additional term in the massless Dirac equation

$$\left(\gamma^\alpha (\partial_\alpha - \Gamma_\alpha) + \xi \gamma^\alpha \gamma^\beta F_{\alpha\beta} \right) \Psi = 0 \quad (2)$$

due to the presence of the anomalous electric moment. It is purpose of this letter to discuss the mechanism of particle creation of electrically neutral massless spin 1/2 particles in the background field (1)

The electromagnetic field, source of the gravitational field with the metric (1) can be written as

$$A_R = \frac{1}{\lambda^{1/4}\chi^{1/2}} \sin t, \quad F_{01} = F_{tR} = \frac{1}{\lambda^{1/4}\chi^{1/2}} \cos t \quad (3)$$

and, the curved Dirac matrices γ^μ can be expressed, in the diagonal (rotating) tetrad gauge, in terms of the flat gamma's $\tilde{\gamma}^\mu$ as follows,

$$\gamma^0 = \lambda^{1/4}\tilde{\gamma}^0, \quad \gamma^1 = \frac{\lambda^{1/4}}{\cos t}\tilde{\gamma}^1, \quad \gamma^2 = \lambda^{1/4}\tilde{\gamma}^2, \quad \gamma^3 = \frac{\lambda^{1/4}}{\sin \vartheta}\tilde{\gamma}^3 \quad (4)$$

then, substituting the gamma matrices γ^μ into the expression for the spinor connections Γ_α

$$\Gamma_\alpha = \frac{1}{4}g_{\mu\lambda} \left[\left(\frac{\partial b_\nu^\beta}{\partial x^\alpha} \right) a_\beta^\lambda - \Gamma_{\nu\alpha}^\lambda \right] s^{\mu\nu} \quad (5)$$

where

$$s^{\mu\nu} = \frac{1}{2}(\gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu) \quad (6)$$

and the matrices b_α^β , a_β^α establish the connection between the curved Dirac matrices γ and the constant Dirac matrices $\tilde{\gamma}$ as follows

$$\gamma_\mu = b_\mu^\alpha \tilde{\gamma}_\alpha, \quad \gamma^\mu = a_\beta^\mu \tilde{\gamma}^\beta \quad (7)$$

we obtain

$$\Gamma_0 = 0, \quad \Gamma_1 = -\frac{1}{2} \sin t \tilde{\gamma}^0 \tilde{\gamma}^1, \quad \Gamma_2 = 0, \quad \Gamma_3 = \frac{1}{2} \cos \vartheta \tilde{\gamma}^2 \tilde{\gamma}^3, \quad (8)$$

Substituting the connections Γ_α into the Dirac equation, and taking into account (3) and (4) we arrive at

$$\left(\tilde{\gamma}^0 \partial_t + \frac{\tilde{\gamma}^1}{\cos t} \partial_R + \tilde{\gamma}^2 \partial_\vartheta + \frac{\tilde{\gamma}^3}{\sin \vartheta} \partial_\varphi + \frac{i\xi}{\chi^{1/2}} \tilde{\gamma}^0 \tilde{\gamma}^1 \right) \Phi = 0 \quad (9)$$

where we have introduced the auxiliary spinor Φ

$$\Psi = (\cos t \sin \vartheta)^{-1/2} \Phi \quad (10)$$

Applying the algebraic method of separation of variables [7, 8], it is straightforward to separate the angular dependence of Φ in (9) from the radial and time ones. Then, after introducing the spinor $\Theta = \tilde{\gamma}^0 \tilde{\gamma}^1 \Phi$ we have that eq. (9) reduces to

$$-i \left(\tilde{\gamma}^2 \partial_{\vartheta} + \frac{\tilde{\gamma}^3}{\sin \vartheta} \partial_{\varphi} \right) \tilde{\gamma}^0 \tilde{\gamma}^1 \Theta = k \Theta \quad (11)$$

$$\left(-\tilde{\gamma}^1 \partial_t - \frac{\tilde{\gamma}^0}{\cos t} \partial_R + \frac{i\xi}{\chi^{1/2}} + ik \right) \Theta = 0 \quad (12)$$

Here, some words about the chirality condition

$$(1 - i\gamma_5)\Phi = 0 \quad (13)$$

are in order, it is not difficult to see that neither the momentum operator in (11) nor eq. (12) commute with $(1 - i\gamma_5)$, then, in order preserve the condition (13) we shall consider as solution of our problem the expression $\Phi_c = (1 + i\gamma_5)\Phi$ which obviously satisfies eq. (13).

Noticing that the operator acting on Θ in eq. (11) is the Brill and Wheeler angular momentum operator [9], we have that the constant of separation k takes integer values $k = \pm 1, \pm 2, \dots$ Using the Dirac matrices representation

$$\gamma^0 = \begin{pmatrix} 0 & i\sigma^2 \\ i\sigma^2 & 0 \end{pmatrix}, \quad \gamma^1 = \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix}, \quad \gamma^2 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \gamma^3 = \begin{pmatrix} 0 & \sigma^3 \\ \sigma^3 & 0 \end{pmatrix} \quad (14)$$

we have eq. (12) can be reduced to the form

$$\left(\frac{d}{dt} - \frac{ik_R}{\cos t} \right) \begin{pmatrix} \Theta_{11} \\ \Theta_{21} \end{pmatrix} - i \left(\frac{\xi}{\chi^{1/2}} + k \right) \begin{pmatrix} \Theta_{22} \\ \Theta_{12} \end{pmatrix} = 0 \quad (15)$$

$$\left(\frac{d}{dt} + \frac{ik_R}{\cos t} \right) \begin{pmatrix} \Theta_{12} \\ \Theta_{22} \end{pmatrix} - i \left(\frac{\xi}{\chi^{1/2}} + k \right) \begin{pmatrix} \Theta_{21} \\ \Theta_{11} \end{pmatrix} = 0 \quad (16)$$

where

$$\Theta = \begin{pmatrix} \Theta_{11} \\ \Theta_{12} \\ \Theta_{21} \\ \Theta_{22} \end{pmatrix} \quad (17)$$

the system of equations takes a more familiar form after making the change of variables

$$t = z + \frac{\pi}{2}, \quad \cos t = -\sin z, \quad (18)$$

then, we look for solutions of the system (15) in the form

$$\Theta_{11} = \sin^{ik_R}(z) \sin\left(\frac{z}{2}\right) I(z), \quad \Theta_{22} = \sin^{ik_R}(z) \cos\left(\frac{z}{2}\right) J(z) \quad (19)$$

where the functions I and J satisfy the system of equations

$$(q+1) \frac{dI}{dq} + (ik_R + \frac{1}{2}) I = -i \left(\frac{\xi}{\chi^{1/2}} + k \right) J \quad (20)$$

$$(q-1) \frac{dJ}{dq} + (ik_R + \frac{1}{2}) J = i \left(\frac{\xi}{\chi^{1/2}} + k \right) I \quad (21)$$

and the new auxiliary variable q reads

$$q = \sin t \quad (22)$$

The solution of the system (20)-(21) can be expressed in terms of hypergeometric functions $F(a, b, \gamma, z)$ as follows:

$$J = {}_2F_1(a, b, ik_R + \frac{1}{2}, (q+1)/2) \quad (23)$$

$$I = -i \frac{\frac{\xi}{\chi^{1/2}} + k}{ik_R + \frac{1}{2}} {}_2F_1(a, b, ik_R + \frac{3}{2}, (q+1)/2) \quad (24)$$

where the parameters a and b read

$$a = ik_R + \frac{1}{2} + \left(\frac{\xi}{\chi^{1/2}} + k \right), \quad b = ik_R + \frac{1}{2} - \left(\frac{\xi}{\chi^{1/2}} + k \right) \quad (25)$$

In order to analyze the phenomenon of particle creation in the background field (1) we proceed to discuss the behavior of the solutions of the Dirac equation in the hypersurfaces $t = \pm\pi/2$, or equivalently when $q = \pm 1$. We can establish the asymptotic behavior of the solutions at the asymptotes by making a comparison with the quasi-classical behavior associated with the Hamilton-Jacobi equation,

$$g^{\alpha\beta} \frac{\partial S}{\partial x^\alpha} \frac{\partial S}{\partial x^\beta} = 0 \quad (26)$$

since the metric $g^{\alpha\beta}$ associated with (1) does not depend on the variables R and φ , we have that the function S can be separated as follows,

$$S = K_R R + k_\varphi \varphi + S_t(t) + S_\vartheta(\vartheta) \quad (27)$$

then, substituting (27) into (26) we arrive at

$$-\left(\frac{dS_t}{dt}\right)^2 + \frac{K_R^2}{\cos^2 t} + \left(\frac{dS_\vartheta}{d\vartheta}\right)^2 + \frac{k_\varphi^2}{\sin^2 \vartheta} = 0 \quad (28)$$

whose solution can be written as

$$\left(\frac{dS_t}{dt}\right) = \pm \sqrt{\frac{K_R^2}{\cos^2 t} + \lambda^2} \quad (29)$$

where λ is a constant of separation associated with the angular dependence of the function S . It is not difficult to see that for $t = \pm\pi/2$ the expression (29) reduces to

$$S_t = \pm K_R \log |\sec t + \tan t| + C, \quad (30)$$

then we have that for $q \rightarrow 1$ $(1-q)^{-ik_R}$ and $(1-q)^{+ik_R}$, and for $q \rightarrow -1$ $(1+q)^{ik_R}$ and $(1+q)^{-ik_R}$ correspond to negative and positive frequency solutions respectively. In order to relate the two different vacua for $q = \pm 1$, we can make use of the relation for the hypergeometric functions $F(\alpha, \beta, \gamma, z)$

$$\begin{aligned} F(\alpha, \beta, \gamma, z) &= \frac{\Gamma(\gamma)\Gamma(\gamma - \alpha - \beta)}{\Gamma(\gamma - \alpha)\Gamma(\gamma - \beta)} F(\alpha, \beta, \alpha + \beta - \gamma + 1, 1 - z) + \\ &+ \frac{\Gamma(\gamma)\Gamma(\alpha + \beta - \gamma)}{\Gamma(\alpha)\Gamma(\beta)} (1 - z)^{\gamma - \alpha - \beta} F(\gamma - \alpha, \gamma - \beta, \gamma - \alpha - \beta + 1, 1 - z) \end{aligned} \quad (31)$$

then the negative frequency mode for $q = -1$ reads

$$\Theta_{(q=-1)-} = c_- \sin^{ik_R}(z) \sin\left(\frac{z}{2}\right) F(a, b, ik_R + \frac{1}{2}, (q+1)/2) \quad (32)$$

whereas the positive energy mode for $q = +1$ takes the form

$$\Theta_{(q=+1)+} = c_+ \sin^{ik_R}(z) \sin\left(\frac{z}{2}\right) F(a, b, ik_R + \frac{1}{2}, (q+1)/2) \quad (33)$$

where c_+ and c_- are constants of normalization. Using the relation (31) and to the fact that

$$\Theta_{(q=-1)-} = \frac{\Gamma(ik_R + \frac{1}{2})\Gamma(-ik_R - \frac{1}{2})}{\Gamma\left(\left(\frac{\xi}{\chi^{1/2}} + k\right)\right)\Gamma\left(-\left(\frac{\xi}{\chi^{1/2}} + k\right)\right)} \frac{c_-}{c_+} \Theta_{(q=+1)+}$$

$$\frac{\Gamma(ik_R + \frac{1}{2})\Gamma(-ik_R - \frac{1}{2})}{\Gamma(ik_R + \frac{1}{2} + (\frac{\xi}{\chi^{1/2}} + k))\Gamma(ik_R + \frac{1}{2} - (\frac{\xi}{\chi^{1/2}} + k))} \frac{c_-}{d_+} \Theta_{(q=+1)-} \quad (34)$$

finally, with the help of the condition $|\alpha|^2 + |\beta|^2 = 1$ for spin $1/2$ particles, we obtain that the density of particles created $|\beta|^2$ is given by the relation

$$|\beta|^2 = \frac{k_R^2 + 1/4}{(\frac{\xi}{\chi^{1/2}} + k)^2} \left| \frac{\Gamma(ik_R + \frac{1}{2})\Gamma(-ik_R - \frac{1}{2})}{\Gamma((\frac{\xi}{\chi^{1/2}} + k))\Gamma(-(\frac{\xi}{\chi^{1/2}} + k))} \right|^2 = \frac{\sin^2(\frac{\xi}{\chi^{1/2}} + k) \pi}{\cosh^2 \pi k_R} \quad (35)$$

The expression (35) shows that the density of created particles has a maximum for $\xi/\chi^{1/2} = N + 1/2$, also we have that, there is not particle creation when $\xi/\chi^{1/2}$ takes integer values. This one is a straightforward consequence of that the eigenvalues k of the angular momentum operator (11) are also integers. Then, for $\xi = 0$ we reobtain the well known result that no particle creation is observed when massless particles couples conformally to the metric.[6] The particular form for $|\beta|^2$ is analogous to the one obtained in analyzing quantum effects in a Robertson Walker metric with an expansion factor $a(\eta) = \varepsilon \cosh^{-1} \eta$, [10, 11] as well as the quantum mechanical problem of tunneling across a barrier of the form $U = -U_0 \cosh^{-2}(\kappa x)$, where for given values of κ the reflection coefficient vanishes.

Acknowledgments

The author wishes to express his indebtedness to the Centre de Physique Théorique for the suitable conditions of work. Also the author wishes to acknowledge to the CONICIT of Venezuela and the Vollmer Foundation for financial support.

References

- [1] L. Parker *Phys. Rev. Lett.* **21** 562 (1968)
- [2] A. A. Grib, S. G. Mamaev, and V. M. Mostepanenko, *Sov. J. Nucl. Phys* **10** 722 (1970)
- [3] B. Bertotti, *Phys Rev.* **116** 1331 (1959)
- [4] I. Robinson, *Bull Acad. Pol. Sci.* **7** 351 (1959)
- [5] V. Sahni , *Phys. Lett. A* **86**, 87 (1981)
- [6] Ya. B. Zeldovich and A. A. Starobinskii *Sov. Phys. JETP* **61** 2161 (1971)
- [7] G. V. Shishkin and V. M. Villalba *J. Math. Phys.* **33** 2093 (1992)
- [8] G. V. Shishkin and V. M. Villalba *J. Math. Phys.* **34** 5037 (1993)
- [9] D. Brill and J. A. Wheeler *Rev. Mod. Phys.* **29** 465 (1957)
- [10] V. M. Villalba and U. Percoco, *Can. J. Phys* **70** 143 (1992)
- [11] A. A. Grib, S. G. Mamaev, and V. M. Mostepanenko. *Quantum vacuum effects in strong fields* (Energoatomizdat 1988)